

IMPLEMENTATION OF THE LYAPUNOV FUNCTIONAL IN DIFFERENCE SCHEMES FOR THE SWIFT-HOHENBERG EQUATION

C. I. Christov

Dept. of Mathematics, University of Louisiana at Lafayette,
Lafayette, LA 70504-1010
christov@louisiana.edu

J. Pontes

Metallurgy and Materials Engineering Department – EE/COPPE/UFRJ PO Box 68505
21945-970 Rio de Janeiro RJ, Brazil
jopontes@ufrj.br

Abstract. Using the operator-splitting method Christov & Pontes (2000, 2001) developed a second-order in time implicit difference scheme for solving the Swift-Hohenberg equation (S-H) which describes pattern formation in Rayleigh-Benard cells. For each time step the scheme involves internal iterations which improve the stability and increase the accuracy with which the Lyapunov functional for S-H is approximated. Different cases of pattern formation were treated and it was shown that the new scheme reaches the stationary pattern several times faster than the previously used first-order in time schemes. In this work we review the main steps concerning the derivation of the second-order in time scheme and prove that the scheme strictly satisfies a discrete approximation of the Lyapunov functional. Results of numerical simulations conducted in four large boxes with two levels of forcings and two different random initial conditions are presented, illustrating that this scheme preserves the non-increasing time-dependent behaviour of the functional. The rate of change of the functional is generally slow save the precipitous downfalls during the time intervals in which the pattern changes qualitatively.

Keywords: Nonlinear Systems, Bénard Convection, Implicit Methods, Finite Difference Methods

1 Introduction

A widely accepted model for the thermal convection in a thin layer of fluid heated from below is the so-called Swift-Hohenberg equation (SH, for brevity) (Swift & Hohenberg, 1977) which is a nonlinear parabolic equation containing fourth-order space-derivatives (generalized diffusion equation). It describes the pattern formation in fluid layers confined between horizontal well-conducting boundaries.

Unlike the models that describe wave propagation on the surface of convective layer, the SH equation possesses a Lyapunov functional which ensures the potential behaviour of the solution. One of the properties of the potential evolution is that only spatial chaos is allowed whereas spatio-temporal chaos is not. However, if one uses a difference scheme which does not faithfully represent the Lyapunov functional one might encounter numerically some non-physical effects. For instance, when the Lyapunov functional is only approximately enforced, then initially the solution approaches one of the attractors in the functional space but eventually leaves the domain of attraction of that particular steady-state solution after being “kicked” from one of the small disturbances the latter arising from the inadequate approximation. Then the solution may keep wandering between several of the attractors exhibiting spatio-temporal chaotic behavior which is not possible for the original model.

Solving SH numerically is a challenge because of the interplay between the higher-order diffusion (higher-order spatial derivatives) and nonlinearity on the one hand, and the presence of a Lyapunov functional – on the other. A computationally efficient difference scheme for SH was developed by Christov et al (1997) where a first-order in time implicit time-stepping is used in the framework of the operator-splitting methods.

There are very few papers in the literature which deal with the problems connected to the numerical implementation of an additional integral constraint on the solution, such as the Lyapunov functional. For dissipative systems (Swift-Hohenberg or complex Landau-Ginzburg equation) an elucidating discussion can be found in the paper by Vinals et al (1991), and the one by Montagne et al (1996). For the case of Lyapunov constraint in a conservative system (such as Nonlinear Schrödinger equation) one is referred

to the work of Ivonin (1999). Though some fine approximations of the additional integral constraint are presented in the above mentioned papers, they are not in fact strictly preserving approximations. We believe, the strict implementation is crucial if one is to deal with more complex models.

In this work we review the main steps concerning the derivation of the second-order in time scheme and prove that the scheme strictly satisfies a discrete approximation of the Lyapunov functional (see also Christov & Pontes 2000, 2001). The scheme employs internal iterations to secure adequate approximation of the nonlinear term. Results of numerical simulations conducted in four large boxes are presented, illustrating that this scheme preserves the non-increasing time-dependent behaviour of the functional.

2 Posing the Problem

Consider a rectangular region $D : \{x \in [0, L_x], y \in [0, L_y]\}$ with boundary ∂D at which different types of b.c. can be imposed. Consider the following Generalized Diffusion Equation (GDE):

$$\frac{\partial u}{\partial t} = -D(\Delta + \kappa^2)^2 + F(u) \equiv -D\Delta\Delta u - 2D\kappa^2\Delta u - D\kappa^4 + F(u). \quad (2.1)$$

For the cubic nonlinearity one has the following potential function

$$F(u) = -\frac{dU(u)}{du} = \varepsilon(x)u - gu^3, \quad U(u) = -\frac{\varepsilon(x)}{2}u^2 + \frac{g}{4}u^4. \quad (2.2)$$

Eq.(2.1) with (2.2) acknowledged is the Swift-Hohenberg equation (SH, for brevity) derived for the Rayleigh-Bénard convection to account for the formation of convective rolls in high Prandtl number fluid layers. The variable $u(x, y, t)$ describes the horizontal planform of the temperature deviation from the conductive profile.

The correct set of lateral b.c. is the one which secures that the evolution of the “energy” $\int v^2$ depends only on its production or dissipation in the bulk, but not on the surface. In other words, in the balance equation for the evolution of energy

$$\begin{aligned} \frac{d}{dt} \frac{1}{2} \int_D v^2 dx dy = & - \oint_{\partial D} v \frac{\partial a_4(x, y, t) \Delta v}{\partial n} dl - \oint_{\partial D} a_4(x, y, t) \Delta v \frac{\partial v}{\partial n} dl - \oint_{\partial D} a_2(x, y, t) v \frac{\partial v}{\partial n} dl - \\ & \int_D a_0(x, y, t) (\Delta v)^2 dx dy - \int_D a_2(x, y, t) (\nabla v)^2 dx dy - \int_D a_0(x, y, t) v^2 dx dy \end{aligned}$$

one has to make the surface integrals vanish. This can happen if one of the following admissible b.c. conditions are imposed

$$v = \frac{\partial v}{\partial n} = 0, \quad v = \Delta v = 0, \quad \frac{\partial v}{\partial n} = \frac{\partial \Delta v}{\partial n} = 0, \quad (x, y) \in \partial D, \quad (2.3)$$

where n stands for the outward normal direction to the boundary ∂D . We call (2.3)₁ and (2.3)₂ “generalized Dirichlet conditions” of the first and second kind, respectively. The condition (2.3)₃ involves only derivatives at the boundary, hence the coinage – “generalized Neumann condition”.

The main feature of (2.1) is that the damping of perturbations occurs via the fourth-order spatial derivatives, while the term with second-order spatial derivatives enhances the perturbations. This allows the occurrence of a linear bifurcation of the solution for certain values of the parameters and/or the size of domain. The nontrivial solutions branch out from a (generally motionless) reference state. Their shapes are the result of the interplay between the complicated linear operator and the nonlinearity. From the perspective of GDE it is clear that a bifurcation can take place only for sufficiently large domains whose size is commensurate with the length scales of the patterns.

SH admits a non-increasing Lyapunov functional

$$\Psi = \int_D \left\{ -\frac{\varepsilon}{2}u^2 + \frac{g}{4}u^4 + \frac{D}{2} [(\Delta u)^2 - 2\kappa^2(\nabla u)^2 + \kappa^4 u^2] \right\} dx dy, \quad (2.4)$$

$$\frac{\partial \Psi}{\partial t} = - \int_D \left(\frac{\partial u}{\partial t} \right)^2 dx dy < 0, \quad (2.5)$$

which rules out the complex temporal or spatio-temporal behavior in the long time (turbulent, oscillatory, or chaotic) and allows formation of steady convective patterns. In their turn these steady patterns can be quite complicated in shape, e.g., spatially chaotic.

Consider the rectangle $x \in [0, L_x]$, $y \in [0, L_y]$ and the Dirichlet b.c. of the first and second kind:

$$u = \frac{\partial u}{\partial x} = 0 \text{ for } x = 0, L_x; \quad u = \frac{\partial u}{\partial y} = 0 \text{ for } y = 0, L_y, \quad (2.6)$$

$$u = \frac{\partial^2 u}{\partial x^2} = 0 \text{ for } x = 0, L_x; \quad u = \frac{\partial^2 u}{\partial y^2} = 0 \text{ for } y = 0, L_y, \quad (2.7)$$

Respectively, the Neumann condition is

$$\frac{\partial u}{\partial x} = \frac{\partial^2 u}{\partial x^2} = 0 \text{ for } x = 0, L_x, \quad \frac{\partial u}{\partial y} = \frac{\partial^2 u}{\partial y^2} = 0 \text{ for } y = 0, L_y, \quad (2.8)$$

There is no restriction to use mixed types of b.c. which are combinations of Dirichlet and Neumann conditions. If the scheme and algorithm perform properly for the “pure” cases (including the Neumann one), then they will do the same for the mixed cases, since any admissible (in the sense of (2.3)) mixture of boundary conditions yields a well posed boundary value problem. For the sake of simplicity we restrict ourselves in the present work to Dirichlet conditions of the first kind.

3 Difference Scheme

3.1 Implicit Time-Stepping

It is not possible to achieve a strict satisfaction of a discrete version of the Lyapunov functional with an explicit scheme which has only two time stages (levels). On the other hand, it is not obvious how the problem can be solved through using multi-level scheme. It is *a-priori* clear that a scheme which is both implicit in time and nonlinear will possess the necessary symmetry to accommodate for the existence of a non-increasing functional. The key to the scheme which satisfies the additional integral constraint (the Lyapunov functional) is the approximation of the nonlinear potential term. A scheme for (2.1) which is implicit and nonlinear reads

$$\frac{u^{n+1} - u^n}{\tau} = D \left[-\frac{\partial^4}{\partial x^4} - \frac{\partial^4}{\partial y^4} - 2k^2 \frac{\partial^2}{\partial x^2} - 2k^2 \frac{\partial^2}{\partial y^2} - 2 \frac{\partial^4}{\partial x^2 \partial y^2} - k^4 \right] \frac{u^{n+1} + u^n}{2} - \frac{U(u^{n+1}) - U(u^n)}{u^{n+1} - u^n}, \quad (3.1)$$

where $U(u)$ stands for the potential of the nonlinear force acting upon the system. For the particular case of quartic potential the nonlinear term adopts the form:

$$-\frac{U(u^{n+1}) - U(u^n)}{u^{n+1} - u^n} = \frac{\varepsilon(x)}{2} [u^{n+1} + u^n] - \frac{g}{4} [(u^{n+1})^3 + (u^{n+1})^2 u^n + u^{n+1} (u^n)^2 + (u^n)^3].$$

3.2 Internal Iterations

The scheme (3.1) is nonlinear and can be solved only by means of iterating the solution within a given time step. Additional benefit from the iterations is that they allow us to alleviate a possible problem connected with the inversion of the linear operators when they are not negative definite. This kind of complications can be expected only when $\max\{L_x, L_y\}k_0 \geq \alpha$, where α can be estimated from the inclusion theorem. As already mentioned, a simple consequence of the non-definiteness of the linear operator is the occurrence of a linear bifurcation of the stationary problem. We tackle this complication by means of an explicit approximation of the second-order terms. Thus the iterative scheme reads

$$\frac{u^{n,k+1} - u^n}{\tau} = \left[-D \frac{\partial^4}{\partial x^4} - D \frac{\partial^4}{\partial y^4} - Dk^4 \right] \frac{u^{n,k+1} + u^n}{2} + \left[-2Dk^2 \frac{\partial^2}{\partial x^2} - 2Dk^2 \frac{\partial^2}{\partial y^2} - 2D \frac{\partial^4}{\partial x^2 \partial y^2} + \varepsilon(x) \right] \frac{u^{n,k} + u^n}{2} - \frac{g}{4} [(u^{n,k})^2 + u^{n,k} u^n + (u^n)^2] u^{n,k+1} - \frac{g}{4} (u^n)^3. \quad (3.2)$$

Here the superscript $(n, k + 1)$ designates the current (“new”) iteration of the unknown set function while the indices (n, k) and (n) distinguish the quantities known from the previous iteration and the previous time step, respectively. The scheme with internal iterations is linear for $u^{n,k+1}$. The internal iterations are conducted until the following criterion is satisfied

$$\frac{\max \|u^{n,K+1} - u^{n,K}\|}{\max \|u^{n,K+1}\|} < \delta, \quad (3.3)$$

for certain $k = K$. Then the last iteration gives the sought function on the new time stage, $u^{n+1} \stackrel{\text{def}}{=} u^{n, K+1}$.

The gist of the concept of internal iterations is that the same time step is repeated until convergence. Since the iterative process begins from an initial condition which is the value of the sought function from the previous time step, the number of internal iterations needed for convergence depends heavily on the magnitude of the time increment τ . For smaller τ the initial condition for the iterations is closer to the sought function and the number of iterations is expected to be small. The trade-off is that a very small τ requires a larger number of time steps which increases the overall number of arithmetic operations per nodal point. Conversely, an inappropriately large τ will bring about a larger number of internal iterations per time step increasing significantly the computational time needed to achieve a single time step dispelling thus the advantage of the larger “strides” (the faster time-stepping). The dependence of the number of internal iterations on τ is nonlinear and leaves a room for optimization. Our numerical experiments show that the calculations are cost effective if the number of internal iterations is in the interval $4 \leq K \leq 16$. This estimate calls for a reduction of the time step when faster processes are treated for which the evolution from a given time stage to the next time stage involves a significant deformation of the field. This means that when faster temporal transients are involved, the usage of larger time steps τ leading to $K \gg 20$ is not justified regardless to the fact that formally speaking the implicit scheme is still stable.

3.3 The Splitting

The inversion of the matrix of Eq. (3.2) is a rather costly procedure even though it is sparse. The 3D case is drastically more expensive. Moreover that the internal iterations require the process to be repeated several times during each time step. Then it is only natural to introduce operator splitting in order to minimize the operations per unit iteration and hence per one time step.

We settle here for the *second Douglas scheme* (Douglas, 1956) (also called “scheme of stabilizing correction” (Yanenko, 1971) which gives the full-time-step approximation for non-commutative operators and is more robust for nonlinear problems than ADI (see Yanenko, 1971) for a review of the splitting schemes and strategies). Another advantage of the stabilizing correction is that for linear problems in 3D it is absolutely stable, while ADI is not. We generalize the Douglas scheme for fourth-order operators and modify it to be second order accurate in time (a Crank-Nicolson type of scheme) as follows

$$\begin{aligned} \frac{\tilde{u} - u^n}{\tau} &= L_{11}^{n,k} \tilde{u} + L_{22}^{n,k} u^n + \frac{1}{2} \left[-D \frac{\partial^4}{\partial x^4} - D \frac{\partial^4}{\partial y^4} - D k^4 + \frac{g}{2} (u^n)^2 \right] u^n \\ &+ \frac{1}{2} [-L_{12} - L_1 - L_2] [u^{n,k} + u^n] \\ \frac{u^{n,k+1} - \tilde{u}}{\tau} &= L_2^{n,k} (u^{n,k+1} - u^n) \end{aligned}$$

where

$$L_{11}^{n,k} \stackrel{\text{def}}{=} -\frac{D}{2} \frac{\partial^4}{\partial x^4} - \frac{D}{4} k^4 + \frac{g}{8} \left[(u^{n,k})^2 + u^{n,k} u^n + (u^n)^2 \right] \quad (3.4)$$

$$L_{22}^{n,k} \stackrel{\text{def}}{=} -\frac{D}{2} \frac{\partial^4}{\partial y^4} - \frac{D}{4} k^4 + \frac{g}{8} \left[(u^{n,k})^2 + u^{n,k} u^n + (u^n)^2 \right] + \frac{\varepsilon(x)}{2} \quad (3.5)$$

$$L_{12} \stackrel{\text{def}}{=} 2D \frac{\partial^4}{\partial x^2 \partial y^2}, \quad L_1 \stackrel{\text{def}}{=} 2D k^2 \frac{\partial^2}{\partial x^2} + \frac{\varepsilon(x)}{2}, \quad L_2 \stackrel{\text{def}}{=} 2D k^2 \frac{\partial^2}{\partial y^2} + \frac{\varepsilon(x)}{2} \quad (3.6)$$

In order to show that the splitting approximates the original implicit scheme we rewrite (3.4) as follows

$$\begin{aligned} [E - \tau L_{11}^{n,k}] \tilde{u} &= [E + \tau L_{22}^{n,k}] u^n + \frac{\tau}{2} [-L_{12} - L_1 - L_2] [u^{n,k} + u^n] \\ &+ \frac{\tau}{2} \left[-\frac{\partial^4}{\partial x^4} - \frac{\partial^4}{\partial y^4} - k^4 + \frac{g}{2} (u^n)^2 \right] u^n, \\ [E - \tau L_{22}^{n,k}] u^{n,k+1} &= \tilde{u} - \tau L_{22}^{n,k} u^n \end{aligned} \quad (3.7)$$

Now we are prepared to eliminate the intermediate variable \tilde{u} . This is done after applying the operator

$[E - \tau L_{11}^{n,k}]$ to the second of equations (3.7) and adding the result to the first one, namely

$$\begin{aligned} [E - \tau L_{11}^{n,k}][E - \tau L_{22}^{n,k}]u^{n,k+1} &= [E + \tau L_{22}^{n,k}]u^n - \tau[E - \tau L_{11}^{n,k}]L_{22}^{n,k}u^n \\ &+ \frac{\tau}{2}[-L_{12} - L_1 - L_2][u^{n,k} + u^n] + \frac{\tau}{2}\left[-\frac{\partial^4}{\partial x^4} - \frac{\partial^4}{\partial y^4} - k^4 + \frac{g}{2}(u^n)^2\right]u^n, \end{aligned}$$

or else,

$$\begin{aligned} [E + \tau^2 L_{11}^{n,k} L_{22}^{n,k}] \frac{u^{n,k+1} - u^n}{\tau} &= (L_{11}^{n,k} + L_{22}^{n,k})u^{n,k+1} + \frac{1}{2}[-L_{12} - L_1 - L_2][u^{n,k} + u^n] \\ &+ \frac{1}{2}\left[-D\frac{\partial^4}{\partial x^4} - D\frac{\partial^4}{\partial y^4} - Dk^4 + \frac{g}{2}(u^n)^2\right]u^n. \end{aligned} \quad (3.8)$$

Upon acknowledging (3.4), (3.5), (3.6), it is readily shown that (3.8) is in fact (3.2) save the *positive definite* operator of norm larger than unity

$$B \equiv E + \tau^2 L_{11}^{n,k} L_{22}^{n,k} = E + O(\tau^2), \quad (3.9)$$

acting upon the time difference $(u^{n+1} - u^n)/\tau$. Acting upon the time difference means that the operator B has no influence on the steady-state result. The fact that $\|B\| > 1$ means that the splitting scheme is more stable than the original implicit scheme. The splitting scheme approximates the desired scheme in full-time steps within the adopted order of approximation $O(\tau^2)$. Thus, employing a splitting does not degrade the temporal approximation of the scheme. In other words, the splitting scheme coincides with the original scheme within the order of approximation of the latter.

4 Implementation of the Lyapunov Functional

After the iterations converge one has $u^{n+1} = u^{n,k+1} = u^{n,k}$ and hence one arrives at a nonlinear scheme in full-time steps which is exactly the difference approximation of scheme (3.1) when in the latter the operators L are replaced by their standard second-order representations in finite differences, Λ (see Christov et al, 1997), namely

$$\frac{u^{n+1} - u^n}{\tau} = (\Lambda_{11} + \Lambda_{22} - \Lambda_1 - \Lambda_2 + 2\Lambda_{12})\frac{u^{n+1} + u^n}{2} + \frac{U(u^{n+1}) - U(u^n)}{u^{n+1} - u^n}, \quad (4.1)$$

where

$$\frac{U(u^{n+1}) - U(u^n)}{u^{n+1} - u^n} = -\frac{\varepsilon}{2}[u^{n+1} + u^n] + \frac{g}{4}[(u^{n+1})^3 + (u^{n+1})^2 u^n + u^{n+1}(u^n)^2 + (u^n)^3].$$

Note that in the left-hand side we neglected the $O(\tau^2)$ contribution to the operator B from (3.9) as asymptotically vanishing with respect to unity.

Upon multiplying eq.(4.1) by $(u_{ij}^{n+1} - u_{ij}^n)/\tau$ and taking the sum over the spatial indices one obtains

$$\begin{aligned} \sum_{i=2}^{M-1} \sum_{j=2}^{N-1} \left[\frac{u_{i,j}^{n+1} - u_{i,j}^n}{\tau} \right]^2 &= \sum_{i=2}^{M-1} \sum_{j=2}^{N-1} \left\{ \frac{\varepsilon}{2\tau} [(u_{i,j}^{n+1})^2 - (u_{i,j}^n)^2] - \frac{g}{4\tau} [(u_{i,j}^{n+1})^4 - (u_{i,j}^n)^4] \right\} \\ &+ \sum_{i=2}^{M-1} \sum_{j=2}^{N-1} \frac{u_{i,j}^{n+1} - u_{i,j}^n}{\tau} [\Lambda_{11} + \Lambda_{22} - \Lambda_1 - \Lambda_2 + 2\Lambda_{12}] \frac{u_{i,j}^{n+1} + u_{i,j}^n}{2} \end{aligned} \quad (4.2)$$

The last term is manipulated using integration (summation) by parts and the discrete form of boundary conditions for u (2.6), (2.7) are acknowledged. We demonstrate the procedure on the difference

operators in x -direction and for Λ_{11} we get

$$\begin{aligned}
\frac{1}{2\tau} \sum_{i=2}^{M-1} u_{ij}^{n+1} \Lambda_{11} u_{i,j}^{n+1} &\stackrel{\text{def}}{=} -\frac{D}{2h_x^4 \tau} \sum_{i=2}^{M-1} u_{ij}^{n+1} [u_{i-2,j}^{n+1} - 4u_{i-1,j}^{n+1} + 6u_{i,j}^{n+1} - 4u_{i+1,j}^{n+1} + u_{i+2,j}^{n+1}] \\
&\equiv -\frac{D}{2h_x^4 \tau} \sum_{i=3}^{M-2} u_{ij}^{n+1} [u_{i-2,j}^{n+1} - 4u_{i-1,j}^{n+1} + 6u_{i,j}^{n+1} - 4u_{i+1,j}^{n+1} + u_{i+2,j}^{n+1}] \\
&= -\frac{D}{2h_x^4 \tau} \sum_{i=2}^{M-1} [2u_{i+1,j}^{n+1} u_{i-1,j}^{n+1} - 4u_{i-1,j}^{n+1} u_{i,j}^{n+1} + 6u_{i,j}^{n+1} u_{i,j}^{n+1} - 4u_{i+1,j}^{n+1} u_{i,j}^{n+1}] \\
&\quad -\frac{D}{2h_x^4 \tau} [-u_{Mj}^{n+1} u_{M-2j}^{n+1} - u_{M-1j}^{n+1} u_{M-3j}^{n+1} - u_{4j}^{n+1} u_{2j}^{n+1} - u_{3j}^{n+1} u_{1j}^{n+1}] \\
&\quad -\frac{D}{2h_x^4 \tau} [4u_{Mj}^{n+1} u_{M-1j}^{n+1} + 4u_{M-1j}^{n+1} u_{M-2j}^{n+1} + 4u_{3j}^{n+1} u_{2j}^{n+1} + 4u_{2j}^{n+1} u_{1j}^{n+1}] \\
&\quad -\frac{D}{2h_x^4 \tau} [-6(u_{M-1j}^{n+1})^2 - 6(u_{2j}^{n+1})^2] = -\frac{D}{2\tau} \sum_{i=2}^{M-1} \left[\frac{u_{i+1,j}^{n+1} - 2u_{i,j}^{n+1} + u_{i-1,j}^{n+1}}{h_x^2} \right]^2
\end{aligned} \tag{4.3}$$

Similarly for Λ_1

$$\begin{aligned}
-\frac{1}{2\tau} \sum_{i=2}^{M-1} u_{ij}^{n+1} \Lambda_1 u_{i,j}^{n+1} &\stackrel{\text{def}}{=} -\frac{2Dk^2}{2h_x^2 \tau} \sum_{i=2}^{M-1} u_{ij}^{n+1} [u_{i-1,j}^{n+1} - 2u_{i,j}^{n+1} + u_{i+1,j}^{n+1}] \\
&= -\frac{2Dk^2}{2h_x^2 \tau} \left\{ \sum_{i=2}^{M-1} \left[u_{i,j}^{n+1} u_{i+1,j}^{n+1} - \frac{1}{2} (u_{i+1,j}^{n+1})^2 - \frac{1}{2} (u_{i,j}^{n+1})^2 \right] + \frac{1}{2} \left[(u_{2j}^{n+1})^2 - (u_{Mj}^{n+1})^2 \right] \right\} \\
&\quad -\frac{2Dk^2}{2h_x^2 \tau} \left\{ \sum_{i=2}^{M-1} \left[u_{ij}^{n+1} u_{i-1,j}^{n+1} - \frac{1}{2} (u_{i,j}^{n+1})^2 - \frac{1}{2} (u_{i-1,j}^{n+1})^2 \right] - \frac{1}{2} \left[(u_{1j}^{n+1})^2 + (u_{M-1j}^{n+1})^2 \right] \right\} \\
&= \frac{2Dk^2}{4\tau} \sum_{i=2}^{M-1} \left\{ \left[\frac{u_{i+1,j}^{n+1} - u_{i,j}^{n+1}}{h_x} \right]^2 + \left[\frac{u_{i,j}^{n+1} - u_{i-1,j}^{n+1}}{h_x} \right]^2 \right\}
\end{aligned} \tag{4.4}$$

Making use of the same technique on can show that

$$\begin{aligned}
\frac{1}{2\tau} \sum_{i=2}^{M-1} u_{ij}^n \Lambda_{11} u_{i,j}^{n+1} - u_{ij}^{n+1} \Lambda_{11} u_{i,j}^n &= 0, & \frac{1}{2\tau} \sum_{j=2}^{N-1} u_{ij}^n \Lambda_{22} u_{i,j}^{n+1} - u_{ij}^{n+1} \Lambda_{22} u_{i,j}^n &= 0, \\
\frac{1}{2\tau} \sum_{i=2}^{M-1} u_{ij}^n \Lambda_1 u_{i,j}^{n+1} - u_{ij}^{n+1} \Lambda_1 u_{i,j}^n &= 0, & \frac{1}{2\tau} \sum_{j=2}^{N-1} u_{ij}^n \Lambda_2 u_{i,j}^{n+1} - u_{ij}^{n+1} \Lambda_2 u_{i,j}^n &= 0.
\end{aligned}$$

In the same vein are treated the terms connected with the y spatial derivative. The approximation of the mixed fourth derivative make use of the above derivations in both x - and y -direction. For the sake of convenience we denote

$$\Phi_{ij} = \frac{u_{i,j-1} - 2u_{ij} + u_{i,j+1}}{h_y^2}.$$

Then

$$\begin{aligned}
\sum_{i=2}^{M-1} u_{ij} \frac{\Phi_{i-1,j} - 2\Phi_{ij} + \Phi_{i+1,j}}{h_x^2} &= \sum_{i=2}^{M-1} \Phi_{ij} \frac{u_{i-1,j} - 2u_{ij} + u_{i+1,j}}{h_x^2} \\
&+ \frac{u_{M-1j} \Phi_{Mj} - u_{1j} \Phi_{2j}}{h_x^2} + \frac{u_{Mj} \Phi_{M-1j} - u_{2j} \Phi_{1j}}{h_x^2} \equiv \sum_{i=2}^{M-1} \Phi_{ij} \frac{u_{i-1,j} - 2u_{ij} + u_{i+1,j}}{h_x^2},
\end{aligned}$$

and hence

$$2 \sum_{i=2}^{M-1} \sum_{j=2}^{N-1} u_{ij} \Lambda_{12} u_{ij} = 2 \sum_{i=2}^{M-1} \sum_{j=2}^{N-1} \frac{u_{i-1,j} - 2u_{ij} + u_{i+1,j}}{h_x^2} \cdot \frac{u_{i,j-1} - 2u_{ij} + u_{i,j+1}}{h_y^2}$$

Now it is readily shown that the right-hand side of (4.2) is the time difference of the Lyapunov functional Ψ

$$\begin{aligned}
\frac{\Psi^{n+1} - \Psi^n}{\tau} &= - \sum_{i=2}^{M-1} \sum_{j=2}^{N-1} \left(\frac{u_{i,j}^{n+1} - u_{i,j}^n}{\tau} \right)^2 \\
\Psi^n &= \sum_{i=2}^{M-1} \sum_{j=2}^{N-1} \left[-\frac{\varepsilon}{2} (u_{i,j}^n)^2 + \frac{g}{4} (u_{i,j}^n)^4 + \frac{Dk^4}{2} (u_{i,j}^n)^2 \right] \\
&\quad - \frac{2Dk^2}{4} \sum_{i=2}^{M-1} \sum_{j=2}^{N-1} \left[\frac{u_{i+1,j}^n - u_{i,j}^n}{h_x} \right]^2 + \left[\frac{u_{i,j}^n - u_{i-1,j}^n}{h_x} \right]^2 + \left[\frac{u_{i,j+1}^n - u_{i,j}^n}{h_y} \right]^2 + \left[\frac{u_{i,j}^n - u_{i,j-1}^n}{h_y} \right]^2 \\
&\quad + \frac{D}{2} \sum_{i=2}^{M-1} \sum_{j=2}^{N-1} \left[\frac{u_{i+1,j}^n - 2u_{i,j}^n + u_{i-1,j}^n}{h_x^2} + \frac{u_{i,j+1}^n - 2u_{i,j}^n + u_{i,j-1}^n}{h_y^2} \right]^2
\end{aligned} \tag{4.5}$$

The last formula presents a $O(\tau^2 + h_x^2 + h_y^2)$ approximation to the original Lyapunov functional (2.5) for the differential equation. The important point about difference version of the Lyapunov functional (4.5) is that it is strictly enforced provided that the internal iterations converge. Its satisfaction does not depend on the truncation error.

5 Numerical Results

The main subject of the present work is to demonstrate the impact of the implementation of the Lyapunov functional on the numerical dynamics. As a featuring example we consider a uniformly loaded system. From the point of view of dynamical systems the behavior of the solution is essentially the same as of the ramped system (Christov et al, 1997) and the only differences are connected with the fact that larger number of solutions with different symmetries can take place for the uniformly loaded system.

Before all we should mention that we checked the spatial and temporal discretizations on a simple case when the whole domain of the flow is occupied by a single convective cell. All mandatory tests involving doubling the temporal and spatial resolutions confirmed the second-order approximation of the scheme and the discrete implementation of the Lyapunov functional. The convergence

of the internal iterations is tested for different values of the tolerance δ defined in eq. (3.3). The quantitative differences between the patterns are negligible for $\delta \leq 10^{-5}$. This allows us to choose $\delta = 10^{-6}$.

In this work we present the results of four numerical simulations conducted in a large box with size $L_x = 50$ and $L_y = 50$ (referred to in what follows as a 50×50 box). The first two simulations were




Figure 1: Results obtained in two simulations run in a 50×50 box with a forcing $\varepsilon = 0.25$ and two different random initial conditions. Steady-state patterns, the time evolution of the associated Lyapunov functional, number of internal iterations accomplished in each time step and time evolution of L_1 are shown from top to bottom. Random initial condition seeds were generated in an IBM SP2 RISC 6000 using the *random* function and seeds 280449 (a) and 123456 (b).

conducted with a forcing $\varepsilon = 0.25$ and two different random initial conditions whereas we adopted $\varepsilon = 0.5$ in the other two and same initial conditions used in the former ones. The results of these four simulations are summarized in Figs. (1) and (2).

For the sake of definiteness and backward compatibility with the works from the literature we select $\kappa_0 = 3.1172$, $g = 12.9$, and $D = 0.015$ which values correspond to a typical Rayleigh-Bénard convection with pattern formation (see, e.g. Swift & Hohenberg, 1977). When the size of the box is $L_x = L_y = 50$, the selected value for k_0 allows roughly 50 convective cells in each direction. A staggered mesh with 302×302 points is used which gives a spatial resolution of approximately 6 points per convective cell (12 points per wavelength). The time step used in all simulations was $\tau = 0.02$.

The random initial condition is constructed by means of a random generator. The value of u in the first point (lower-left corner) is calculated with the random generator for a given initial seed and then the rest of the grid points are filled row-wise using the previous point as a seed for the next point. Finally, the initial field is renormalized to $[-1,1]$. Thus, the pattern can be referred by the number of the initial seed. We assess the rate of evolution of the pattern during the simulation by monitoring the relative L_1 norm defined as

$$L_1 = \frac{1}{\tau} \frac{\sum_{i,j} |u_{i,j}^{n+1} - u_{i,j}^n|}{\sum_{i,j} |u^{n+1}|}, \quad (5.1)$$

which roughly corresponds to the ratio between the spatial average of the modulus of time derivative u_t and the spatial average of the modulus of the function itself.

The calculations begin from the random initial condition and proceeded until $L_1 \leq 5 \times 10^{-7}$, when it can be assumed that the motion is virtually steady. In all simulations the Lyapunov functional $\Psi(t)$ and L_1 rapidly decrease at the very beginning of the evolution. In this stage most of the spectral content of the random initial condition is filtered and only wavelengths close to κ_0 survive. This is the stage in which the structure is formed and the amplitude of the pattern saturates. The pattern is characterized by a large number of defects and short correlation lengths. The second stage is characterized by elimination of defects, resulting in longer correlation lengths. The *phase* of the patterns evolve with minor changes in the average amplitude. The Lyapunov functional decreases at a much lower rate and L_1 norm evolves slower. However, the evolution of L_1 is not monotonical and accelerates when defects collapse. This behaviour is precisely captured by the sudden increases of the norm $L_1(t)$ and abrupt decreases in the Lyapunov functional.

We monitored also the number of internal iterations. In the time intervals of slow evolution, the number of internal iterations is around 1-3. As expected, this number increases in the time intervals of rapid evolution (the same intervals in which the L_1 norm increases).

Manipulating the initial condition we succeeded to obtain a plethora of solutions as illustrated in




Figure 2: Results obtained in two simulations run in a 50×50 box with a forcing $\varepsilon = 0.50$ and two different random initial conditions. Steady-state patterns, the time evolution of the associated Lyapunov functional, number of internal iterations accomplished in each time step and time evolution of L_1 are shown from top to bottom. Random initial condition seeds were generated in an IBM SP2 RISC 6000 using the *random* function and seeds 280449 (a) and 123456 (b).

Figs. (1) and (2). The effect of increasing the forcing ε applied to the system is also shown in the figures: higher forcings lead to a higher density of defects and shorter correlation lengths due to the fact that a wider band of wavelengths becomes linearly unstable.

6 Conclusions

In the present work we review and applied the operator-splitting difference scheme proposed by Christov and Pontes (2000, 2001), for the numerical solution of nonlinear diffusion equations containing fourth-order space-derivatives. The scheme is of second-order approximation both in time and space and does not contain artificial dispersion, hence the disturbances are quickly attenuated. It is fully implicit owing to the use of internal iterations. The main characteristic of the scheme is that a discrete version of the Lyapunov functional which holds for the original differential equation is strictly implemented.

The performance of the scheme is demonstrated for the evolution of the solution of the Swift-Hohenberg (SH) equation in four simulations conducted in large boxes with two forcing levels and two different random initial conditions. The SH equation models the Rayleigh-Bénard convection in a horizontal layer. The strict implementation of the Lyapunov functional with its non-increasing behaviour is clearly demonstrated. The numerically obtained Lyapunov functional decreases more rapidly when the pattern undergoes qualitative changes (when the solution enters the immediate vicinity of the respective attractor). In all cases the evolution ended up in a spatially chaotic pattern. No temporal chaos was possible due to the strict satisfaction of the Lyapunov functional.

The proposed difference scheme can serve as a model for constructing approximations to nonlinear physical system when additional constraints, such as Lyapunov functional, are imposed on the solution.

7 Acknowledgements

The work of C.I.C. is supported in part by Grant LEQSF(1999-2002)-RD-A-49 from the Louisiana Board of Regents. J. P. acknowledges financial support from CNPq (Brazil) through grant No. 452399/00-9(NV). The simulations were performed on the IBM SP2 RS-6000 of the NACAD-UFRJ Computer Center at the Federal University of Rio de Janeiro, Brazil.

References

- CHRISTOV, C. I., AND PONTES, J. A second-order scheme for evolution equations modelling Bénard convection. In *Proceedings of the 8th Brazilian Congress of Thermal Engineering and Sciences* (Brazil, 2000). Paper S06P01 (CD).
- CHRISTOV, C. I., AND PONTES, J. Numerical scheme for Swift-Hohenberg equation with strict implementation of the Lyapunov functional. *Mathematical and Computer Modelling* ((To appear)).
- CHRISTOV, C. I., PONTES, J., WALGRAEF, D., AND VELARDE, M. G. Implicit time splitting for fourth-order parabolic equations. *Comput. Methods Appl. Mech. Engrg* 148 (1997), 209–224.
- DOUGLAS, J., AND RACHFORD, H. H. On the numerical solution of heat conduction problems in two and three space variables. *Trans. Amer. Math. Soc.* 82 (1956), 421–439.
- IVONIN, I. A., PAVLENKO, V. P., AND PERSSON, H. Self-consistent turbulence in the two-dimensional nonlinear Schrodinger equation with a repulsive potential. *Phys. Rev. E* 60 (1999), 492–499.
- MONTAGNE, R., HERNADEZ-GARCIA, E., AND SAN-MIGUEL, M. Numerical study of a Lyapunov functional for the complex Ginzburg-Landau equation. *Physica D* 96 (1996), 44–65.
- SWIFT, J., AND HOHENBERG, P. C. Hydrodynamic fluctuations at the convective instability. *Phys. Rev. A* 15 (1977), 319–328.
- VINALS, J., HERNADES-GARCIA, E., SAN MIGUEL, M., AND TORAL, R. Numerical study of the dynamic aspects of pattern selection in the stochastic Swift-Hohenberg equation, in one dimension. *Phys. Rev. A* 44 (1991), 1123–1133.
- YANENKO, N. N. *Method of Fractional Steps*. Gordon and Breach, NY, 1971.